

<http://www.liebertonline.com/doi/abs/10.1089/pho.2008.2350>

Comparison Between Single-Diode Low-Level Laser Therapy (LLLT) and LED Multi-Diode (Cluster) Therapy (LEDT) Applications Before High-Intensity Exercise

To cite this article:

Ernesto Cesar Pinto Leal Junior, Rodrigo Álvaro Brandão Lopes-Martins, Bruno Manfredini Baroni, Thiago De Marchi, Rafael Paolo Rossi, Douglas Grosselli, Rafael Abeche Generosi, Vanessa de Godoi, Maira Basso, José Luis Mancalossi, Jan Magnus Bjordal. Photomedicine and Laser Surgery. -Not available-, ahead of print.
doi:10.1089/pho.2008.2350.

Online Ahead of Print: March 20, 2009

Ernesto Cesar Pinto Leal Junior, M.Sc.,^{1,2,3}
Rodrigo Álvaro Brandão Lopes-Martins, Ph.D.,⁴
Bruno Manfredini Baroni, P.T.,^{1,2}
Thiago De Marchi,^{1,5}
Rafael Paolo Rossi,^{1,5}
Douglas Grosselli,^{1,7}
Rafael Abeche Generosi, P.E.,^{1,2,6}
Vanessa de Godoi,^{1,5}
Maira Basso,^{1,5}
José Luis Mancalossi,^{1,5} and
Jan Magnus Bjordal, Ph.D.^{3,8}

¹Laboratory of Human Movement, University of Caxias do Sul, Caxias do Sul, RS, Brazil.

²Sports Medicine Institute, University of Caxias do Sul, Caxias do Sul, RS, Brazil.

³Section for Physiotherapy Science, Institute for Public Health and Primary Health Care, University of Bergen, Bergen, Norway.

⁴Laboratory of Pharmacology and Phototherapy of Inflammation, Department of Pharmacology, Institute of Biomedical Sciences, University of São Paulo, São Paulo, SP, Brazil.

⁵Faculty of Physiotherapy, University of Caxias do Sul, Caxias do Sul, RS, Brazil.

⁶Federal University of Rio Grande do Sul, Porto Alegre, RS, Brazil.

⁷Faculty of Physical Education, University of Caxias do Sul, Caxias do Sul, RS, Brazil.

⁸Bergen University College, Institute for Physical Therapy, Bergen, Norway.

Address reprint requests to:

Ernesto Cesar Pinto Leal Junior, M.Sc.

Laboratory of Human Movement

Sports Medicine Institute (IME)

University of Caxias do Sul

Rua Francisco Getúlio Vargas, 1130

Caxias do Sul, RS 95070-560,

Brazil

E-mail:

Abstract

Background Data and Objective: There is anecdotal evidence that low-level laser therapy (LLLT) may affect the development of muscular fatigue, minor muscle damage, and recovery after heavy exercises. Although manufacturers claim that cluster probes (LEDT) maybe more effective than single-diode lasers in clinical settings, there is a lack of head-to-head comparisons in controlled trials. This study was designed to compare the effect of single-diode LLLT and cluster LEDT before heavy exercise. **Materials and Methods:** This was a randomized, placebo-controlled, double-blind cross-over study. Young male volleyball players (n=8) were enrolled and asked to perform three Wingate cycle tests after 4×30 sec LLLT or LEDT pretreatment of the rectus femoris muscle with either (1) an active LEDT cluster-probe (660/850 nm, 10/30 mW), (2) a placebo cluster-probe with no output, and (3) a single-diode 810-nm 200-mW laser. **Results:** The active LEDT group had significantly decreased post-exercise creatine kinase (CK) levels (-18.88 ± 41.48 U/L), compared to the placebo cluster group (26.88 ± 15.18 U/L) ($p < 0.05$) and the active single-diode laser group (43.38 ± 32.90 U/L) ($p < 0.01$). None of the pre-exercise LLLT or LEDT protocols enhanced performance on the Wingate tests or reduced post-exercise blood lactate levels. However, a non-significant tendency toward lower post-exercise blood lactate levels in the treated groups should be explored further. **Conclusion:** In this experimental set-up, only the active LEDT probe decreased post-exercise CK levels after the Wingate cycle test. Neither performance nor blood lactate levels were significantly affected by this protocol of pre-exercise LEDT or LLLT.